

SUPPLEMENTARY MATERIAL TO: Naidoo et al. S Afr J Sci. 2020;116(5/6), Art. #7693, 8 pages.

HOW TO CITE:

Naidoo T, Rajkaran A, Sershen. Impacts of plastic debris on biota and implications for human health: A South African perspective [supplementary material]. S Afr J Sci. 2020;116(5/6), Art. #7693, 5 pages. <https://doi.org/10.17159/sajs.2020/7693/suppl>

Table 1: Summary of published data on marine plastic debris ingestion in shark, fish, turtle, bird and mammal species from South Africa and surrounding islands

Taxon	Species	Feeding	Type	Size (mm)	Region	Location	Environment	% Incidence	Items per individual (± s.d.)	Source
Sharks										
Sand tiger shark	<i>Carcharias taurus</i>	Carnivorous	PP, PE, F	plastic bags	KZN		C/O	0.04% of 2268		1
Copper shark	<i>Carcharias brachyurus</i>	Carnivorous	PP, PE	plastic sheets	KZN		C/O	0.06% of 1404		1
Scalloped hammerhead	<i>Sphyrna lewini</i>	Carnivorous	PP, PE		KZN		C/O	0.1% of 1916		1
Great hammerhead	<i>Sphyrna mokarran</i>	Carnivorous	–		KZN		C/O	0 of 177		1
Smooth hammerhead	<i>Sphyrna zygaena</i>	Carnivorous	PP, PE		KZN		C/O	0.17% of 1154		1
Dusky shark	<i>Carcharias obscurus</i>	Carnivorous	PP, PE		KZN		C/O	0.15% of 2741		1
Black tip shark	<i>Carcharias limbatus</i>	Carnivorous	PP, PE		KZN		C/O	0.22% of 1785		1
Bull shark	<i>Carcharias leucas</i>	Carnivorous	PP, PE		KZN		C/O	0.61% of 661		1
Shortfin mako shark	<i>Isurus oxyrinchus</i>	Carnivorous	PP, PE		KZN		C/O	0.87% of 231		1
Tiger shark	<i>Galeocerdo cuvier</i>	Carnivorous	PP, PE		KZN		C/O	7.52% of 505		1
Java shark	<i>Carcharhinus amboinensis</i>	Carnivorous	–		KZN		C/O	0 of 222		1
Spinner shark	<i>Carcharhinus brevipinna</i>	Carnivorous	–		KZN		C/O	0 of 1699		1
Sandbar shark	<i>Carcharhinus plumbeus</i>	Carnivorous	–		KZN		C/O	0 of 379		1
Great white shark	<i>Carcharodon carcharias</i>	Carnivorous			KZN		C/O	0.38% of 524		1

Taxon	Species	Feeding	Type	Size (mm)	Region	Location	Environment	% Incidence	Items per individual (± s.d.)	Source
Bony fish										
Razorbelly	<i>Hilsa kelee</i>	Zoobenthivorous	Fb	<1	KZN	Durban Harbour	C	9 ± 3.5	2	
Silver silago	<i>Sillago sihama</i>	Zoobenthivorous	Fb	<1	KZN		C	6 ± 5.2	2	
Pursemouth	<i>Gerres filamentosus</i>	Zoobenthivorous	Fb	<1	KZN	Durban Harbour	C	8 ± 4.7	2	
Glassfish	<i>Ambassis dussumieri</i>	Zoobenthivorous	Fb,F	0.76 ± 0.77	KZN	Durban Harbour	C	69% of 29	0.93 ± 0.75	3
Tilapia	<i>Oreochromis mossambicus</i>	Zoobenthivorous	Fb,F	0.64 ± 0.46	KZN	St Lucia Estuary	C	38% of 29	0.41 ± 0.57	3
Thornfish	<i>Terapon jarbua</i>	Zoobenthivorous	Fb,F	0.94 ± 0.81	KZN	Umgeni Estuary	C	48% of 29	0.66 ± 0.81	3
Juvenile mullet	<i>Mugil</i> sp.	Microphytobenthos	Fb,F	1.09 ± 0.77	KZN	Durban Harbour	C	55% of 29	1.00 ± 1.46	3
Sub-adult mullet	<i>Mugil cephalus</i>	Microphytobenthos	F,Fb,Fl,PS	0.2–15	KZN	Durban Harbour	C	73% of 70	3.8 ± 4.7	4
Turtles										
Loggerhead	<i>Caretta caretta</i>	Benthic carnivorous	F,P,Fl	<30	WC	Cape Town	C	60% of 40	7.7 ± 13.9	5
Loggerhead	<i>Caretta caretta</i>	Jellyfish			WC	Cape Agulhas	C	11% of 37		6
Birds										
Blue petrel	<i>Halobaena caerulea</i>	crustaceans, fish	P,F,Fb, Fm	1–20	SA/SO		B/C/O	92% of 74		7
Great shearwater	<i>Ardenna gravis</i>	fish, squid, crustaceans	P,F,Fb, Fm	1–20	SA/SO		B/C/O	90% of 50		7
White-faced storm petrel	<i>Pelagodroma marina</i>	crustaceans, fish	P,F,Fb, Fm	1–20	SA/SO		B/C/O	88% of 24		7
Pintado petrel	<i>Daption capense</i>	crustaceans, fish	P,F,Fb, Fm	1–20	SA/SO		B/C/O	83% of 18		7
Wilson's storm petrel	<i>Oceanites oceanicus</i>	fish, squid, crustaceans	P,F,Fb, Fm	1–20	SA/SO		B/C/O	75% of 4		7
Thin-billed prion	<i>Pachyptila belcheri</i>	zooplankton and fish	P,F,Fb, Fm	1–20	SA/SO		B/C/O	69% of 32		7

Taxon	Species	Feeding	Type	Size (mm)	Region	Location	Environment	% Incidence	Items per individual (± s.d.)	Source
Antarctic priori	<i>Pachyptila desolata</i>	crustaceans, fish, squid	P,F,Fb, Fm	1–20	SA/SO		B/C/O	59% of 88	7	
White-chinned petrel	<i>Procellaria acquinocialis</i>	crustaceans, fish	P,F,Fb, Fm	1–20	SA/SO		B/C/O	57% of 201	7	
Salvin's prion	<i>Pachyptila salvini</i>	crustaceans, fish	P,F,Fb, Fm	1–20	SA/SO		B/C/O	52% of 31	7	
Sooty shearwater	<i>Ardenna grisea</i>	fish, squid, krill	P,F,Fb, Fm	1–20	SA/SO		B/C/O	51% of 63	7	
Grey phalarope	<i>Phalaropus fulicarius</i>	zooplankton	P,F,Fb, Fm	1–20	SA/SO		B/C/O	50% of 2	7	
Arctic skua	<i>Stercorarius parasiticus</i>	fish, birds	P,F,Fb, Fm	1–20	SA/SO		B/C/O	50% of 2	7	
Cory's shearwater	<i>Calonectris diomedea</i>	fish, squid, crustaceans	P,F,Fb, Fm	1–20	SA/SO		B/C/O	43% of 7	7	
White-bellied storm petrel	<i>Fregetta sp.</i>	squid, crustaceans	P,F,Fb, Fm	1–20	SA/SO		B/C/O	38% of 13	7	
Grey-backed storm petrel	<i>Garrodia nereis</i>	squid, crustaceans	P,F,Fb, Fm	1–20	SA/SO		B/C/O	33% of 12	7	
Broad-billed prion	<i>Pachyptila vittata</i>	crustaceans, squid	P,F,Fb, Fm	1–20	SA/SO		B/C/O	30% of 137	7	
Kerguelen petrel	<i>Aphrodroma brevirostris</i>	squid, crustaceans, small fish	P,F,Fb, Fm	1–20	SA/SO		B/C/O	24% of 63	7	
Subantarctic skua	<i>Catharacta antarctica</i>	seabirds, molluscs, fish	P,F,Fb, Fm	1–20	SA/SO		B/C/O	23% of 494	7	
Soft-plumaged petrel	<i>Pterodroma mollis</i>	squid, crustaceans	P,F,Fb, Fm	1–20	SA/SO		B/C/O	21% of 29	7	
Kelp gull	<i>Larus dominicanus</i>	squid, crustaceans	P,F,Fb, Fm	1–20	SA/SO		B/C/O	13% of 52	7	
Black-browed albatross	<i>Thalassarche melanophrys</i>	fish, crustaceans	P,F,Fb, Fm	1–20	SA/SO		B/C/O	11% of 18	7	
Antarctic fulmar	<i>Fulmarus glacioloides</i>	crustaceans, squid, fish	P,F,Fb, Fm	1–20	SA/SO		B/C/O	11% of 27	7	
Atlantic petrel	<i>Pterodroma incerta</i>	squid, crustaceans	P,F,Fb, Fm	1–20	SA/SO		B/C/O	10% of 20	7	
Great-winged petrel	<i>Pterodroma macroptera</i>	squid, crustaceans	P,F,Fb, Fm	1–20	SA/SO		B/C/O	8% of 13	7	
Southern giant petrel	<i>Macronectes giganteus</i>	fish, crustaceans	P,F,Fb, Fm	1–20	SA/SO		B/C/O	7% of 123	7	

Taxon	Species	Feeding	Type	Size (mm)	Region	Location	Environment	% Incidence	Items per individual (\pm s.d.)	Source
Northern giant petrel	<i>Macronectes halli</i>	krill, fish, squid	P,F,Fb, Fm	1–20	SA/SO		B/C/O	7% of 42		7
Antarctic petrel	<i>Thalassoica antarctica</i>	krill, fish, squid	P,F,Fb, Fm	1–20	SA/SO		B/C/O	7% of 30		7
Subantarctic shearwater	<i>Puffinus elegans</i>	squid, fish	P,F,Fb, Fm	1–20	SA/SO		B/C/O	7% of 15		7
Snow petrel	<i>Pagodroma nivea</i>	fish, molluscs	P,F,Fb, Fm	1–20	SA/SO		B/C/O	5% of 22		7
Wandering albatross	<i>Diomedea exulans</i>	octopus, squid, cuttlefish	P,F,Fb, Fm	1–20	SA/SO		B/C/O	4% of 156		7
Atlantic yellow-nosed albatross	<i>Thalassarche chlororhynchos</i>	fish, crustaceans	P,F,Fb, Fm	1–20	SA/SO		B/C/O	2% of 87		7
Common diving petrel	<i>Pelecanoides urinatrix</i>	crustaceans	P,F,Fb, Fm	1–20	SA/SO		B/C/O	2% of 53		7
Southern rockhopper penguin	<i>Eudyptes chrysocome</i>	fish, squid	P,F,Fb, Fm	1–20	SA/SO		B/C/O	1% of 177		7
Grey-headed albatross	<i>Thalassarche chrysostoma</i>	crustaceans, fish, squid	P,F,Fb, Fm	1–20	SA/SO		B/C/O	1% of 170		7
Sooty albatross	<i>Phoebetria fusca</i>	squids, crustaceans	P,F,Fb, Fm	1–20	SA/SO		B/C/O	1% of 73		7
Bank cormorant	<i>Phalacrocorax neglectus</i>	fish, crustaceans	P,F,Fb, Fm	1–20	SA/SO		B/C/O	1% of 167		7
Mammals										
Subantarctic fur seal	<i>Arctocephalus tropicalis</i>	Myctophid fish							0	8
Antarctic fur seal	<i>Arctocephalus gazella</i>	Myctophid fish							0	8

Type: PP, polypropylene; PE, polyethylene; F, fragments; Fb, fibres; Fl, film; PS, polystyrene; P, pellets; Fm, foam

Region: KZN, KwaZulu-Natal; SA, South Africa; SO, Southern Ocean; WC, Western Cape

Environment: C, coastal; O, oceanic; B, beach

References

1. Cliff G, Dudley SF, Ryan PG, Singleton N. Large sharks and plastic debris in KwaZulu-Natal, South Africa. Mar Freshw Res. 2002;53(2):575–581.
2. Naidoo T, Goordiyal K, Glassom D. Are nitric acid (HNO_3) digestions efficient in isolating microplastics from juvenile fish? Water Air Soil Pollut. 2017;228(12):1–11. <https://doi.org/10.1007/s11270-017-3654-4>
3. Naidoo T, Sershen, Thompson RC, Rajkaran A. Quantification and characterisation of microplastics ingested by selected juvenile fish species associated with mangroves in KwaZulu-Natal, South Africa. Environ Pollut. 2020;257, Art. #113635. <https://doi.org/10.1016/j.envpol.2019.113635>
4. Naidoo T, Glassom D, Smit AJ. Plastic ingestion by estuarine mullet, *Mugil cephalus* (L., 1758), in an urban harbour, KwaZulu-Natal, South Africa. Afr J Mar Sci. 2016;38(1):145–149. <https://doi.org/10.2989/1814232x.2016.1159616>
5. Ryan PG, Cole G, Spiby K, Nel R, Osborne A, Perold V. Impacts of plastic ingestion on post-hatchling loggerhead turtles off South Africa. Mar Pollut Bull. 2016;107(1):155–160. <https://doi.org/10.1016/j.marpolbul.2016.04.005>
6. Hughes GR. The sea turtles of South-east Africa. 1. Status, morphology and distributions. Oceanographic Research Institute investigative report 35. Durban: South African Association for Marine Biological Research; 1974.
7. Ryan PG. The incidence and characteristics of plastic particles ingested by seabirds. Mar Environ Res. 1987;23(3):175–206. [https://doi.org/10.1016/0141-1136\(87\)90028-6](https://doi.org/10.1016/0141-1136(87)90028-6)
8. Ryan PG, De Bruyn PN, Bester MN. Regional differences in plastic ingestion among Southern Ocean fur seals and albatrosses. Mar Pollut Bull. 2016;104(1–2):207–10. <https://doi.org/10.1016/j.marpolbul.2016.01.032>